n.d.

LIBRARY
RECEIVED

\*\*

U.S. Department of Agridulture

# VEGETABLE and FLOWERING Annual Seeds

FOR

Cold, Hot and Rainy Seasons and also Miscellaneous Seeds



PROCURABLE AT THE

Empress Botanical Gardens,

POONA.

India

A SALE AND THE SALE OF THE SAL

#### MEMORANDUM

All the Vegetable and Flower Seeds enumerated in this list have been obtained direct from well-known firms in Europe and America. They are strongly recommended as being of the best quality procurable. The seeds are true to name, are tested every three months, and the present list contains only such kinds as a long experience has proved to thrive best here. The seeds are issued on the same conditions on which they have been received from European firms and the undersigned gives no guarantee, express or implied, as to description, quality, productiveness, or any other matter of any seeds, and will not be in any way responsible for the crop, there being too many causes, known and unknown, which prevent good seeds from germinating.

Unless orders are received to the contrary, the nearest available varieties will be substituted for those out of stock.

All indents should be addressed to the Superintendent.

Indentors are specially requested to write name and address clearly and to give forwarding instructions where necessary.

Indents not accompanied with a remittance cover the cost of seeds, packing charges and postage in full, will be complied with by value payable system.

The Postmaster-General has ruled that seeds supplied on payment are not 'bona fide' trade samples of no saleable value and therefore cannot be despatched by sample or pattern post.

The rate of postage per value-payable and per ordinary parcel post is the same, viz., 3 annas per lb. Registration fee will be charged Re. 0-2-0 per parcel of any value.

The maximum weight carried at above rates is I1 pounds.

In all instances packing charges will be kept as near as, possible to the cost price of the material employed; but special packing charges will be made on large collections of small seeds and on bulky seeds of low value.

The seed godown is open on all days, except Sundays and holidays, between the hours 8 to 11 A.M., and 3 to 6 P.M. By order,

N. M. BHAGAWAT.

Empress Botanical Gardens. Poona. range and remaining to the beautiful and rest estimate

Superintendent

# NOTES ON SEED GROWING

#### Why Seeds Fail

Vegetables or Flowers of good quality cannot be grown without first sowing genuine seeds; it is exceedingly annoying after preparing, sowing and cultivating, to find that instead of having spent time and money profitably, you have lost heavily in obtaining a spurious article. When purchasing be satisfied that the party from whom you obtain supplies is reliable, or that such a house has a good reputation, and expect to pay a fair price for what you obtain. There are plenty of cheap and worthless seeds, as well as any other article in the market and it would be always advisable to guard against them.

Some cultivators, through forgetfulness of the fact that the products of a garden, being natives of various soils and climates, require peculiar management, sow the seeds in the ground at improper seasons.

Some seeds, such as those of Beans, Cabbage, Lettuce, Radish, Turnip &c., being from their nature apt to vegetate quickly, are often destroyed while germinating through variableness of the weather; others are liable to be devoured by insects in forty-eight hours after they are sown, and before a plant is seen above the ground unless a suitable remedy is applied in time to destroy the vermin.

Other species, such as Beet, Carrots, Celery, Leek, Onion, Parsley, Spinach, &c., being naturally of hardy growth, take in favourable seasons from two to three or four weeks to vegetate, are apt to perish through incrustation of the soil, or other untoward and unaccountable circumstances which cannot al-

ways be controlled.

Failures often occur through seeds being deposited too deeply in the ground, or left too near the surface. Sometimes for the want of a proper quantity of seed in a given spot solitary plants will perish they not having sufficient strength to open the pores of the earth, and very frequently injudicious management in over manuring and improperly preparing the soil causes defeat. Other failures of success are, for instance, the shooting or prematurely going to seed of certain vegetables, as Lettuce, Endive &c.; the reason in most cases is to be found in sowing too thick and in leaving the plants which are to be transplanted too long in the seed bed, or in thinning them out too late if they are to be left on the spot; often also in checking the growth by keeping too dry, or through ravages of insects, and frequently by a spell of hot weather, which forces up the growth instead of swelling the heart.

The seedsman's fair reputation is often unjustly defamed, through the failure of the seeds, when the fault is to be attributed entirely to one or other of the above causes.

# REMEDIES AGAINST FAILURE OF SEEDS

First—A proper and constant degree of moisture without being soaked with water. This is secured by making the surface of freshly dug soil so fine that the smallest seeds may cone in immediate contact on all sides with the particles from which they are to absorb the required moisture and the pressing of the soil over the seeds so firmly with the feet or the back of a spade that the degree of moisture may remain as nearly as possible the same until the plants are up.

Second—A proper degree of heat, secured by sowing each variety of seed when the average temperature of the locality is that most favourable for its germination. This may be learned

from a careful study of the practice of the locality.

Third—Covering the seed to such a depth that while it is preserved at a uniform degree of heat and moisture, the necessary air can readily reach it, and the tiny stem pushes the forming leaves into the light and air. This depth will vary with different seeds and conditions of the soil and can be learned only from practical experience. In general, seeds of the size of the Turnip should be covered with half an inch of earth pressed down, while Beans may be two or three and Peas two to four inches deep.

Fourth—Such condition of soil that the ascending stem can easily penetrate it, and the young roots speedily find suitable food. We can usually secure this by thorough preparation of the ground, and taking care never to sow fine seeds when the ground is wet. Occasionally a heavy or long continued rain followed by a bright Sun will so bake and crust the surface that it is impossible for the young plant to find its way through it, or a few days of strong wind will so dry the surface that no seed will germinate. In such cases the only remedy is to plant again.

Fifth—Remove every weed as soon as it appears, if well done while the plants are young much after-work will be saved

Cultivation of the ground.—The soil as it naturally occurs in most gardens contains sufficient material to favour the growth of plants.

To obtain cultivated plants in perfection, two operations require to be attended to in order to ensure a successful result; the proper cultivation of the ground or tilth and the supply of

manurial ingredients. After the earth is in a state of fine division by being dug over several times a liberal supply of decayed manure and leaf-mould should be dug in and after the soil has been suitably arranged and raked smooth, it can be slightly watered in anticipation of planting or sowing.

Manure.—A general manure for flower beds is supplied by the stable and cow shed manure, leaf mould, ashes and all kinds of refuse usually stored in pits in compounds. For gross feeders, (when growing vigorously only and not when they are at rest) such as Cannas, Chrysanthemums, most tuberous and bulbous plants, a weekly or bi-weekly application of liquid manure is necessary.

The method of preparation is simple, consisting merely of stirring farmyard manure into a vessel of water—a wooden tub by preference. The mixture should be stirred occasionally and replenished as it weakens. The clear fluid only, (free of sediment) should be used and it is advisable to dilute it in the water-can before pouring it on the soil.

Sowing the seeds:—Seeds can be sown either in beds protected against the extreme heat or light of the sun or in shallow boxes or pots in an open verandah. The latter course is advisable in the rainy weather and always with seeds of rare or valuable plants, as a watch can be more easily maintained against the attacks of insects, birds, and other possible enemies, and the owner if need be, can also more directly supervise the watering and other delicate operations.

Care must be taken that the soil is of extreme fineness throughout and that the drainage is good. In a bed this can be arranged for by raising its level; holes to act as out-lets should be bored in the bottoms of the pots or boxes.

In sowing, the seeds should be scattered evenly, and in the case of small seeds, a covering of fine soil not more than about a quarter of an inch in depth should be sifted over them and lightly patted firm with the palm of the hand or with some flat tool, as a slight firmness on the surface assists the germinating seed to rid itself of its outer skin or coat. Larger seeds should be inserted about an inch beneath the surface. In watering, great care should be taken not to flood the seed bed; to keep the soil moist is all that is necessary, and it is safer to use watering can with a very fine rose.

To judge when seedlings are in the most favourable stage for transplanting is a matter of practice, but it is generally time to do so when six leaves have formed. Some seeds such as those of Mignonette, Poppy, Candytuft, etc., should be sown where they are to remain as these plants do not bear transplanting well.

Transplanting of Seedlings:—This should be done in the evening by preference. In very sunny weather it is as well to protect the transplanted seedlings by shading them with leafy twigs for a day or two.

Before commencing operations the seed bed should be watered if it be not already sufficiently moist, and then opened at one side or end to below the levels of the tips of the roots. Each individual seedling should be gently raised upwards with a narrow flat piece of wood and an attempt should be made at the same time to bring up a small ball of earth in which the roots are imbedded. To plant the seedling a hole should be made in the ground deep and wide enough to take the roots without bending them. Holding the base of the stem between thumb and forefinger just at the surface of the bed, the earth can be filled in round the roots and finally pressed down slightly to retain the plant in position. Utmost gentleness in handling the seedlings is necessary, as their young growing parts are so tender and delicate. The intervals apart at which seedlings should be placed depends upon the ultimate spread of each kind.

Treatment of plants during growth:—Many plants such as Dahlias, Chrysanthemums, Hollyhocks, Cosmos etc., require to be supported by stakes while growing or when full grown to prevent their being blown down by wind or by the weight of their flowers.

The soil in beds should not be allowed to become too dry and at the same time, care should be taken not to allow it to remain in a sodden state. Stirring of the soil and the accompanying extermination of weeds at frequent intervals should not be neglected.

Some of the important Monsoon annuals.—Aster, Balsam, Candytuft, Celosia, Cosmos, Dahlia, Double Daisy, Gaillardia, Hollyhocks, Lobelia, Marigold, Nasturtium, Phlox, Pinks, Salvias, Torenia, Verbena, and Zinnias.

Bulbous Plants for the Rainy Season:—Achimenes, Tuberous-rooted Begonias, Caladiums, Crinums, Dahlias, Gesneras, Gladiolus, Gloriosa Superba, Gloxinias, Pancratiums and Zephyranthes (Crocus).

# VEGETABLE SEEDS

Note—The cultural directions are given in a general way to guide amateurs, and may have to be changed to suit soil and climate. Every attempt is made to stock all the varieties of seeds mentioned but some may not be in stock and selections of an alternative variety may be made.

#### VEGETABLE SEEDS

Artichoke—SOW in August, broadcast in a bed of rich loose soil, and when the seedlings are large enough to handle, plant out in the open, three to four feet either way.

Price per oz. Rs. 2—0

Artichoke purple, or purple Artichoke French green.


Asparagus (A delicate vegetable)—SOW in August to December.—SOIL dug to two feet deep, manured heavily, and dressed with salt.—WATER keep soil moist and cool.

Warm exposure. Tender shoots ready third year.

Price per oz. As. 8

Asparagus purple, early, giant | Asparagus, Erfurt giant white.

Argenteuil



Dwarf French Bean Wax Dale


Beans French—SOW 6 inches apart at intervals of a fortnight. in the rainy season or from the end of cold season, in furrows about two feet apart—SOIL must be deep and rich with manure—WATER should be given in large quantities once a week in dry weather.

Beans, Dwarf, Kidney.	. 1	Rs.	a.
do Canadian Wonder true select	per lb.	2	0
do. Best of all	do.	2	Ü
do. Yellow podded wax, golden butter	do.	2	0
Beans, Runner			
do. Scarlet and scarlet mammoth	do.	2	0
do. Painted lady	do.	1	8
do. Lima or Double	do.	2	8
Beans Broad Windsor green long pod	do.	1	12
do. Seville, Earliest	do.	1	12
beans French, Kidney, acclimatized	do.	0	12



## Beet Egyptian dark red turnip rooted

Beet—SOW any time throughout the year, 10 in a sapart in lines 15 inches apart, on the sides of ridges—SOIL, rich well-tilled. Thin out and gently transplant when the seedlings are small, leaving only one in each place.—WATER freely in dry weather.



# Price per oz. As. 6

Beet, Crimson globe

do. Eclipse, very early.

Egyptian earliest dark red. do.

do. Erfurt long black red.

do. Long black, cameroon.

do. Perpetual or Spinach.

White silver Seakale, Improved do. Spinach.

Swiss chard yellow, cutting. do.

Borecole.—SOW between June and September, in beds of fine soil.-PROTECT from insects that abound in July, by ashes and watering overhead.—PLANT OUT on the sides of ridges 21 feet apart when about 6 leaves have grown.

Price per oz. As. 8.

Borecole or Kale, Dwarf green-curled, extra selected.


tion a second of and

do. Thousand-headed.

Best Bow we had been be Beet, long black Cameroon

Broccoli Learnington ... per oz. 2

Culture: The same as for Cauliflower. Does not flower in the plains.



Rs. As.


Brussels Sprouts, Reading
Exhibition

per oz. 1 -8

do. Dwarf Improved

do. 1 0





Cabbage largest Schweinfurt

#### Cabbage Savoy earliest Ulm.

Cabbage—SOW about the middle of the rainy season in a bed of fine soil,—TRANSPLANT when five or six leaves have formed large sorts 30 to 36 inches apart each way, small curly sorts 24 inches apart—SOIL must be very rich—WATER enough to keep the soil moist. Dig in a fresh supply of strong manure when the leaves begin to curve inwards. Use Hellebore powder or Tobacco soap against caterpillars and green fly.



Cabbage, Burnswick.

Drumhead (Fottler's)


•		Ks.	As.,
Cabbage, Blood Red, Erfurt, early	per	oz. 1	8
do. Brunswick (Fottler's) large white	do.	1	ξ.
do. Drumhead or Scotch	do.	1	. 8
do. Early Flat Dutch (Premium)	do.	1	8
do. Schweinfurt, Quintal largest white	do.	1	8
do, Sugar loaf	do.	1	8
do. Savoy (Sugar loaf, Ironhead, Vienna Ulm			
vars.)	do.	2	0
do. American Florida Header	do.	- 2	0
do. do. Glory of Enkhuizen	do.	$^2$	0
do. do. Prize medal Drumhead	do.	2	0
do. do. Copenhagen Market, early	do.	2	. 0



Capsicum.—SOW in a seed bed, if rainfall is light about beginning of rainy season, at end, if heavy. Transplant when the seed-lings are about 6 inches high—SOIL rich, well-tilted and manured. WATER freely.

Capsicum monstrous large sweet.

	*				7	As.
(	apsic	um Elephant's trunk red	***	per 1/4	oz. packet	6
	do.	Spanish Mammoth, red			do.	6
	do.	Bull nose or large bell			do.	6
	do.	Monstrous or grossum, red	•••		do.	6
	do.	do. Golden yellow			do.	6
,	do.	Chinese giant, scarlet	•••		do.	- 6
	do.	Ruby King	* * *		do,	. б



Carrot, Scarlet Amsterdam


Carrot—SOW any time between beginning of rains and end of cold season in drills one foot apart.—SOIL any fair garden.—WATER during dry weather.

		£	ls.
Carrot, Scarlet Amsterdam	<i>₹</i> ₩•	per oz.	8
	- <b>T</b>	do.	8
do. do. Oxheart	1.00	do.	8
do. Altringham, long scarlet	www. Table 1	do.	8
do. long red Surrey	***	do.	3
do. Country, pink, hardy	4.7 h	đó.	3



Cauliflower, Snowball.

Cauliflower—SOW about middle of the rainy season, on a bed of very fine rich soil—TRANSPLANT when five or six leaves have formed, 30 inches apart—SOIL should be heavily manured and drawn round the plants as they grow, and constantly stirred and hoed, WATER enough to keep soil moist, give liquid manure frequently.



Cauliflower V. A. Giant

per oz.

Cauliflower, Erfurt
very dwarf small
leaved, snowball 5 0

do. Algiers, extra
selected ... 4 0

do. Danish large
early ... 3 0

do. Veitch's Autumnal giant ... 2 8

do. Early London
or Dutch early 2 8

Rs



Celery White Solid

Celery—SOW about beginning of rainy season, in a bed of fine rich soil—TRANS-PLANT into trenches, lines 2 feet apart plants 1 foot apart—SOIL very rich, free working and well-drained.—WATER very freely, giving liquid manure once a week—BLANCH by tying up in drain pipes and straw when well-grown. An open sunny position is best.

#### Price per oz. Re. 1-4-0.

Celery, white-ribbed, large dwarf solid.


do. ' white, Pascal Giant.

Cress.—SOW in a bed of very rich soil, thickly in lines, a small quantity every three days, throughout the year. SHADE from bright sunshine—Water daily.

Cress, extra curled ... ... Per oz. As. 2 do. Water, improved, broad leaved per 1/8 oz. packet As. 4

Cucumber.—SOW any time from beginning of hot weather to end of the rainy season in rich well-tilled soil.—TRAIN on a trellis or stakes. Some varieties spread on the ground—WATER enough to keep the soil moist.

	As.	
Cucumber, white Parisian long ridge	per $\frac{1}{4}$ oz. 4	
do. Green, long Athens		
do. the dold Excelsion to the final house of	do. 4	
do. Half long green, white-spined	do. 4	
do. Japanese climbing	do. 4	
do. Country	do. 4	



Brinjal, Purple Giant.

Egg Plant.—Aubergine or Brinjal. SOW if rain-fall is light
any time during the rainy season; if heavy sow when rain is
nearly over, on a bed of rich
soil— TRANSPLANT when
five or six leaves have formed
2 to 3 feet apart. WATER
when soil is dry, giving liquid
manure frequently.

•	Aā.
Egg-plant, purple long per ½ oz. do. Long white giant d	packet 4
do. Long white giant d	lo. 4
do. New York giant round purple	lo. 6
do. Black Beauty, enormous d	lo. 6
do Oblong striped pink	o. 3
Endive, moss curled extra fine and the per of	z. 6
do. Broad leaved white	o. 6
<b>6</b> 11	,

Culture.—The same as for Lettu ce.

Gooseberry Cape or Tiparee.—SOW on a seed bed at the beginning of the rains.—TRANSPLANT when seedlings are about 8 inches high, in furrows 3 to 4 feet apart either way—Fill the furrows by drawing soil from the ridges, as the plants grow. Crop ready November onwards.

As.

Gooseberry cape

per ½ oz. packet 6



Knol Kohl.—Or Kohl Rabi—SOW from beginning of rains till end of cold season, except where rain-fall is very heavy, m beds of fine soil—TRANSPLANT when about six leaves are made—SOIL should be well worked and manured.

Price per oz. Re. 1

Knol Kohl, green early earliest Erfurt

do. green or white late giant. do. do. Goliath.

Knol Kohl early white Vienna short leaved.

do. purple late giant, Goliath

Leek—SOW in a well-manured seed bed—TRANSPLANT when about 9 inches high, in trenches a foot deep, at one foot apart, gradually earth up to blanch. A gross feeder.

Price per oz. Re. 1-4-0

Leek, broad leaved flag or London.

do. winter Erfurt

Leek, Carentan giant, hardy.

do. do. Musselburgh.



Lettuce, Butter Cup

Lettuce—SOW if rainfall is light fortnightly between the beginning of the rains and the end of the cold season, if heavy, from the end of the rains TRANSPLANT a foot apart when 5 or 6 leaves are thrown out. SOIL, must be rich but not lately manured. WATER, keep the soil moist. Tying round with plantain shreds induces "heading",



Lettuce, Tom Thumb.

		Rs.	A.
Lettuce, Cabbage, Butter Cup	per oz.	1	0
do. do. Paris Sugar	do.	0	12
do. do. Drumhead or Malta	<b>d</b> o.	0	12
Lettuce, Cos. Paris white, extra			
selected.	<b>d</b> o.	0	12
do. Trianon early white self-folding	do.	0	12
do. Dwarf yellow self-folding	do.	0	12
do. Red or Brown	do.	0	12

Lettuce for picking or gathering (These form heads more than once after each cutting)

WATER, freely-SHADE slightly.

Mangelwurzel-This is a root crop grown in Europe like Sugar The cultivation is the same as for ordinary Beet Beet. root (for salads). But Mangels are always grown for cattle food. The roots are gross feeders and when fully grown weigh 5 to 10 lbs.

BB. 5 Mangelwurzel long red ner oz. do. 0 5 Sugar Beets



Melon Scarlet



As.

12

per oz.

Melon yellow netted

Melon.—SOW about middle of cold season.—SOIL, very sandy and heavily manured.—WATER very freely.

#### Price 1 oz. As. 4

Melon, Netted and other sorts | Melon, Water mixture mixed. many sorts. do. Malta red fleshed. do. do. Wonderful sugar.

Mustard.—SOW every three days from beginning of rains to end of cold weather. SOIL must be rich and fine.

		• •				As.
Mustard, finest large white	1	***			0	5
do. finest brown	•••	4.6		do.	0	5



Onion, Madeira Globe

Onion.—For Green Onions.— SOW fortnightly throughout rainy and cold season. For Ripe Onions SOW September to November, on a bed of rich soil-TRANS-PLANT, when 9 inches high 4 to 6 inches apart,

Price per oz. Re. 1-8-0.

Onion, red globe acclimatized (As. 8)

white globe do. (As. 8) do.

Lisbon late giant white. do.

do. Italian Mammoth white.

do. Prizetaker vellow. Onion, Welsh white or English

do. Zittan giant yellow.

Tripoli, giant Rocca do. (12 As.)

do. Madeira Globe.

Parsley.—SOW any time from beginning of rainy to middle of hot season, on bed where it is to remain.—SOIL should be deep and sandy. WATER—Frequently while small. SHADE—Afternoon, in hot weather.

Price per oz. As. 8

Parsley, Beauty of the Parterre | Parsley, Hybrid moss-curled.

do. Fern leaved.

Triple-curled dwarf extra.

do. Double curled.

do. Dwarf Perfection.

Parsnip.—SOW any time between the end of rains and middle of cold season in ridges 2 feet apart. SOIL rich, welltilled. WATER—Freely in dry weather.

Price per oz. As. 8

Parsnip, Long smooth. do. Sutton's Student. Parsnip, round early.



Pea-SOW if rainfall is light, from beginning of rain to middle of cold sea-2 to 3 feet son in tows SUPPORT distant. With stakes when seedlings are 6 inches high SOIL.—Should be manured for previous crop. WA-TER-Enough to keep soil constantly moist. BE-WARE — of mildew. draught encourages, moisture prevents it.

Price per lb. Rs. 2.

Pea American Wonder Marrow.

do. Champion of England do.

do. Knight's dwarf Marrow (Sugar).


do. Excelsior, Suttons.

Pea World's Record.

do. Telephone

do. Triumph

do. Country (As. 12 per lb.)



Hundred Weight Yellow.

Pumpkin—SOW in rich friable soil during the rains. The seed should be sown (2 or 3) a few inches apart, in patches 10 feet one from the other.

Price per 1 oz. As. 3.

Pumpkin,	large	hundred
	weig]	ht green
do.		Scarlet
do.		Yellow
do		White



Radish red round

Radish.—SOW any time during the rainy, cold and hot seasons in lines 12 inches apart, or scatter broad-cast in loose free working soil.—SOIL rich well tilled. Do not transplant,—WATER freely in dry weather.



Radish, flat red



Radish, long red



Radish slender

# Price per oz. As. 8.

	Radish, long scarlet, white-
do. do vellow	tipped
do. do. white.	do. long, rose, white-tipped
do. scarlet, Olive shaped.	do. long, white Vienna cr lady finger.
do. do. French break-	do. long, white country
fast.	(As. 3.)

Rhubarb.—American Giant

per oz. Re. 1-0

Rosel

per oz. As. 12

Salsify, or vegetable Oyster, Sow in drills. The roots have the flavour of Oysters and are cooked per oz. Re. 1-8

The state of the state of the state of

Spinach.—SOW if rainfall is light, from beginning of rainy to middle of cold season in lines 18 inches apart, and at 12 inches distance. SOIL must be rich and friable. WATER enough to keep soil constantly moist. SHADE Afternoon.

#### Price per oz. As. 6


Spinach, monstrous Viroflay. | Spinach, Victoria or Market favourite.

do. Bloomsdale. | do. acclimatized As. 4.

Strawberry.—Large fruited sorts mixed Price per 1 oz. As 10

Sugar Corn or Table Maize.—This produces daintiest of cobs, with very tender grains, full of milky sweet juice. Will grow best in the rains, though sowings on small patches can be made till the middle of the cold season. Will succeed best in back yards where the soil is usually rich. Price per pound Rs. 2-8.

Sugar Corn Crosbys; early, First of all. Wrighled, deinen



Tomato, King Humbert



Tomato, Large red.

Tomato.—SOW, if rainfall is under 50 inches from beginning of rainy to middle of cold season; fresh—sowing monthly

—PLANT OUT—When a few inches high in lines five feet apart, and four feet between plants. SOIL—Must be well drained. WATER—Freely to keep soil moist and liquid manure often. Provide stout stakes to keep the fruit off the ground, when about a foot high.

Price Re. 1-8 per oz.

Tomato, Mikado Scarlet.

do. Trophy, large, red, smooth.

do. Peachblow.


do. King Humbert.

Tomato, Perfection.

do. Ponderosa.

do. Large red (As. 12 per

do. Magnum Bonum.



Turnip purple top Munich

from the beginning of rainy season to end of cold season, in lines 18 inches apart and thin out to 12 inches distant. SOIL.—Must be rich and often hoed. Only one plant in each hole should be kept.

Price per oz. As. 8.


Turnip, early Dutch.

do. Early Milan red top.

do. purple top Munich.

do. White snow ball.

do. Golden yellow.



Yogetable Marrow scalloped Turk's Cap and the control of

March August and October. SOIL Rich and thoroughly drained. WATER profusely during dry weather.

Price per oz. Re 1

Vegetable Marrow or squash, custard marrow, white and golden.

do. scalloped Vars, mixed.

do. long white bush mar.

do. do. yellow.

do, cream.

do, best sorts mixed.

#### POT HERBS

#### Packet 4 Annas

- Basil Sweet—A favourite pot herb. Easily grown in the rains from seeds sown in pots or ground.
- Lavender—SOW after rains, in light rich bed or pans and transplant when three inches high into small pots or beds.
- Marjoram—SOW in the beginning of November in pans in very light soil, and transplant when big enough to handle.
- Peppermint—SOW at the beginning of the cold weather in a seed bed and then transplant a foot apart.
- Rosemary—SOW September to November, in pans in very light soil, and transplant when large enough to handle.
- Sage—A favourite pot herb.—SOW between August and December.—SOIL any fine garden—WATER, enough to keep the soil moist. SHADE afternoon.
- Tarragon—SOW after rains in very light soil in pans and transplant 5 or 6 inches apart.
- Thyme—A favourite pot herb—SOW between August and December.—SOIL, deep sandy—WATER, enough to keep soil moist.
- Wormwood—A Sweet scented herb sown at the end of the rains. When dry, is used for scenting clothes:
- NOTE.—The proper mint is different from Peppermint and is propagated by roots only. Sold at 8 As., a bundle of cuttings.

#### COUNTRY VEGETABLE SEEDS.

(Price per ounce, except where mentioned otherwise.)

Per oz. Rs. As. Bhendi or Okra (Hibiscus esculentus) Lady's finger Bhopla or Kadu (Cucurbita maxim,) 0 12 Brinjals Bengali (Solanum melong ana) Baigan 0 12 Dorli ( do. . . ) do. Chakwat (Chenopodium album) Spinach Chandanbatwa (Artiplex hortensis) do. ... 0 2 Chawlai (Amarantus polygamus) do. 2 0 Chawli (Vigna cattiang- chawli bean) 2 Chawdhari or Goa bean (Posphocarpus tetragonolobus) (per doz. seeds) ... 0 Chillis (caspicum furtescens) 1 Chooka (Rumex vesicarius) Spinach 0 8 Dilpasant (Citrullus vulgaris) Tinda gourd 0 12 Dudhi (Lagenaria vulgaris) Bottle gourd long or round... 0 3 Gawar (Cyampsis psoraloides) Bean 4 Ghosali or Ghee turai (Luffa aegyptica) gourd 0 4 Kardai (Carthamus tinctorius) Saff ower 0 1 Karela green (Momordica charantia) Bitter gourd 0 8 do. white 1 0 Kashiphal (Cucurbita Pepo) ... 0 6 Kohala (Benicasa cerifera) Gourd melon 0 12 Kothmir (Coriandum Sativum) Indian parsley 1 Maize (Indian corn) Makai Boota (per lb.) 0 8 Math (Amarantus blitum) Spinach  $\mathbf{2}$ Methi (Trigonella foenumgroecum) (per lb.) 1 0 Mogri (Serpentine radish pods) 0 8 Mula (Raphanus Sativus) 3 0 Padwal (Trichosanthes anguinia) Snake gourd 1 4 Palak (Spinacia Oleracea) 0 2 Pokla (Amarantus blitum. var) Spinach 0 2 Rajgira ( do. salicifolius ) 2 Shepu or Sowa (Peucedanum graveolens) 2 Turai or Dodka (Luffa acutangula) Ridge gourd 4 Walhor or Ghevda varieties (Doliche's lablab var) 6 0 Walpapdi do. do. 2

# FLOWER SEEDS.

-:0:--

Note.—The price of most flower seeds is annas four per packet; where it is higher, it is given against that particular kind.

**Acroclinium** (e y e r - lasting flowers) roseum.

do. (ever-lasting flowers) album.


for many days. Grows about

This is an annual that succeeds both in the rains as well as in the cold weather. The flowers last like paper flowers 15 inches high.



#### Ageratum

Ageratum mexicanum blue and white. This annual produces a mass of flowers during the cold weather. It will also succeed in the rains, if light. Height about one foot.



Alyssum.

Alyssum, Sweet, Little Gem. (fragrant).

do. rostratum dwarf yellow.

Grows about 6 inches high. This makes a good girdle of white to a round bed of dwarf, coloured annuals. Does best during the cold season.



Amarantus bicolor ruber.

Amarantus.— Annual with very bright showy coloured leaves.—SOW during rainy or cold season—SOIL very rich.—WATER freely, during dry weather.

Amarantus, tricolor (Jacob's coat) showy.

do. tricolor Splendens very beautiful foliage.

do. bicolor ruber

do. Salicifolius (fountain plant)

do. Caudatus (Love-lies bleeding).



Amarantus Caudatus.



Amarantus Tricolor.



#### Antirrhinum Queen Victoria.

Antirrhinum, or Snapdragon. A very showy annual.—SOW, during rainy or cold season.—SOIL, rich friable— WATER, enough to keep the soil moist.

Antirrhinum, tall vars, finest | Antirrhinum. Semi-dwarf vars mixed.

do. vars, large flowering finest mixed. As. 6.

best mixed.

do. Queen Victoria white (As. 6.)

Aquilegia single vars, mixed. Aquilegia double vars, mixed. (As. 8) This does not flower in the tropical climate, but requires a temperate atmosphere.

Arctotis—Bluish white large, daisylike flowers which keep open from 9 a.m. to 5 p.m. They open by lamp-light in vases.

Arnebia Cornuta.—A charming cold season annual growing to a height of 2 feet and bearing spotted yellow flowers.


Asparagus—A hardy green house climber, feathery green foliage which looks graceful in Bouquets and Table Decorations. Sow seeds in pots. Transplant singly, Germination slow.

per pkt. Re. I: do. plumosus do. A . 8 Sprengeri do.



Aster Dwarf Pyramidal Bouquet,

Aster—COLOUR, white, lilac, crimson and intermediate shades. HEIGHT, 6 inches to 1½ feet—SEASON rainy and cold. SOW, at beginning of rainy season if rainfall is under 50 inches. if greater sow at end of rainy season in fine mould, in pots or boxes. TRANSPLANT, when two inches high.—WATER daily during dry weather.



Aster,—Boltze's Dwarf Bouquet, Finest mixed.

do. Tall, Quilled German, Finest mixed.

do. Mignon, Finest mixed.

do. Victoria Giant, Finest mixed.

do. Paeony-flowered, Finest mix d.

do. Ostrich Plume, Salmon Rose.

do. Ostrich Plume, Crimson.

do. Ostrich Plume, Gloriosa, ruby coloured.

do. Ostrich Plume, White.

do. Ostrich Plume, Rose.

do. Ostrich Plume, Finest mixed.

do. Giant Comet, Finest mixed.



Aster Comet.



Balsam Camelia.




Balsam Rose Flowered.

Balsam—COLOUR, white, crimson and intermediate shades— SOW, between May and August every 30 days to keep up a succession.—SOIL, must be very rich and friable. Water daily, giving liquid manure, frequently.

Balsam double rose flowered. splendid mixture.

double rose flowered white, rose, scarlet, crimson, separate colours (Per pkt. As. 6.)



do. Carnation striped finest mixed (As. 6.)

do. finest mixture of rose flowered and spotted.



Bellis Perennis or double daisy.—SOW in pans and cover lightly with leaf-mould. TRANSPLANT in beds about 6 inches apart.

Bellis perennis (Double Daisy) mixed very fine quality. (Per pkt. As. 8.)

Cacalia Coccinea.—An annual (rainy and cold season) with very showy thistle-like red flowers.

do. lutea (aurantiaca) orange yellow.

Calceolaria grandiflora, Tigered finest mixed (Re. 1-8) Doer well in Hills only: in the plains if the cold season is long.

Calendula officinalis fl. ple, grandiflora. Golden yellow. Grows anywhere in the rains.

Calliopsis Bicolor—A showy annual—SOW, during rainy and cold season—SOIL, any good—WATER daily.

Calliopsis, Bicolor tinctoria | Callopsis mixed dwarf sorts. free flowering.

red.

astrosarguinea do. do.

do. maxima new large fl.

do. Drummondi.

Campanula medium (Canterbury Bell) single mixed double do.

Canterbury Bells usually take two years to flower unless specially forced. Sow when heavy rains are over, where intended to bloom or in pots.



Candytuft—COLOUR, putest white, also purple and other shades—HEIGHT, 6 o 10 inches.—SOW July to December in light rich soil where intended to grow.

Candytuft, Rose
Cardinal
Candytuft, Emperor
large.

Candytuft, Umballata purpurea crimson.

Candytuft, White rocket.

Canna.—Hybrida.—COLOUR, Primrose, red, crimson, &c.—
HEIGHT,—1½ to 4½ feet.—SEASON, hot and rainy.
SOW April to August. Previously soak the seed in
manure water for three days, or rub lightly on a sharp
file.—SOIL very rich—WATER, freely during dry weather, less during cold season.

Canna, many sorts mixed.

do. dark leaved varieties splendid mixed.

do. newest large flowering varieties mixed (As. 8.)

Carnation,—Lovely plants for pots or beds.—SOW, in July or August in pans in light sandy soil. Plant out in beds or pots when about two inches.—WATER daily in dry weather.

Carnation, Border, early, double flowering, dwarf, Vienna,

	•	per pkt. Re. 1
do.	do. white	do. do.
do.	double marguerite finest mixed	do. do.
do.	do. dark red	do. do.
do.	perpetual or tree, very fine mixture	do. do.



Celosia Peramydalis Thompsoni.



Cock's comb.

Celosia Cristata. Cock's comb — COLOUR, crimson, yellow &c .--HEIGHT, 6 to 18 inches—SEASON, rainy and cold-SOW at beginning of rainy season if rainfall is under 50 inches, if greater -SOW about end of rainv season in smull pots, plant out when seedlings are 3 inches high -WATER, daily during dry weather.

Celosia Cristata (cock's comb) nana (dwarf) very fine quality mixed.

do. do. Empress. (As. 12)

do. do. finest mixed selected quality.

do. Pyramidalis Thompsoni (feathery) (As. 8.)

do. do. plumosa splendid mixed.


Centaurea, Cornflower or Blue bottle—SOW, from beginning of rainy to end of cold weather, in small pots or boxes and plant out when the seedlings are about 3 inches high in beds about 6 inches apart—Should be out in the Sun—WATER so as to keep the bed cool.

Centaurea cyanus minor, mixed.

do. do. Emperor William, blue.

do, do dwarf Victoria blue.





Chrysanthemum Carinatum.

Chrysanthemum Inodorum.

Chrysanthemum.—COLOUR, sulphur, white, &c.—HEIGHT 1 to 3 feet.—SOW, about beginning of rainy season. SOIL—any fair garden soil.—WATER, daily. The various Japanese and other scented Chrysanthemums are grown by suckers.

Chrysanthenum Carinatum mixed.

do. Frutescens (Paris Daisy).

do. Leucanthemum grandiflorum (Ox Eye Daisy).

do. Leucanthemum Maximum (Shasta Daisy).

do. Coronarium double mixed. do. Inodorum (Bridal Robe).

Cineraria—For foliage and flowering plants—SOW seed in September in pans or boxes and prick out seedlings when 5 or 6 leaves have been made. A cold weather annual C. Maritima vars. make splendid edgings for beds or as centre pieces for green beds.

Cineraria, Maritima acanthifolia

do. do. candidissima

do. do. Grandiflora mixed Per pkt. Re. I.

Clarkia—The usual instructions for sowing and planting as for cold weather annual. Flowers good for cutting and effective in beds.

Clarkia fine mixed single.

do. do. double.

Cleome Speciosa—Sow in the place in which wanted to bloom or sow in seed beds and transplant, at the beginning of the rains; a bed of them makes a mass of purple if planted about 18 inches apart; or plant at the back of low beds.



Coleus.

- Coleus. SOW seeds in pots and transplant singly when about 2-3 inches high. Various colours from seeds. Cuttings root freely, start cuttings from April.
- do. Finest quality splendid mixture (As. 12.)
- do. laciniated leaved mixed. (Re. 1.)
- do. large leaved mixed (Rs. 1-8.)

Convolvulus—COLOUR, white, rose, purple and intermediate shades.—HEIGHT, 10 feet climbing.—SEASON raing —SOW at the beginning of the rainy season, where it is intended to bloom—SOIL, deep, friable, rich—WATER freely.



Convolvulus tricolor.

- Convolvulus tricolor or minor best mixed (for beds.)
  - do. major best mixed
    (Morning Glory, for trellis or arches.)
  - do. Japanese Morning Glory (for beds).

Coreopsis—COLOUR, bright yellow—HEIGHT, 1 foot,—SEASON, rainy and cold—SOW, if rainfall is under 50 inches from the beginning of the rainy to the middle of the cold season. SOHL, any fair garden—WATER, enough to keep soil moist.

Coreopsis grandificra, large golden yellow do. fl. pleno, new double (As. 8.)

Cosmos—(Cosmea bipinnata) COLOUR, white, pink, purple, orange.—SOIL any garden, out in the sun—SOW from June to August, in a seed bed and plant in beds when seedlings are about 4 inches high, about 12 inches apart each way.

—WATER enough to keep soil moist.

Cosmea	bipinnata alba (white)	Cosmea	
1.	7 / 1 1 .		Klondyke (golden
do.	do. rosea (pink).		yellow). do. mammotin mix-
do.	do, purpurea (pur-	do.	ed.
		do.	do. double mixed
	ple.)		Re. 1 per pkt.

Cuphea—fine mixed. Sow in the rains. Good for cutting, Various Colours. A hardy annual 10 to 15 inches high. Cyclamen—Persicum Re. 1-8 pretty upturned flowers.

Dahlia—A very showy annual with flowers of bright colours—SOW, April to August.—SOIL very rich.—WATER daily till the flowers are over. The roots can be stored in the cold weather to be planted in May next.

Dahlia double mixed very fine quality	Per pkt. As. 8	
do. do. Cactus varieties do. Giant Perfection	do. 8	;
do. Giant Perfection	do. 8	5
do. Collarette	. do 8	ţ.
do. Lucifer		
do. Tom Thumb (dwarf)		
do. single vars splendid mixed		
do. single Cactus new vars		

Delphinium elatum hybridum, double and single finest mixed (As. 6). Grow only in the cold, season like Larkspurs, Charming flowers. Succeed best in the hills.

Dhatura—COLOUR, white, cream, &c.—SOW any time during the year—SOIL, will grow in any soil.—WATER, sufficiently to keep the soil moist.

Dhatura, double finest mixed.



Dianthus
Indian Pink Double



Dianthus Barbatus Sweet William

Dianthus.—Indian or Chinese Pink.—SOW between June and December—SOIL, rich friable.—WATER daily. Hardy and showy annual, Good for cutting.

Dianthus Barbatus (Sweet William) mixed, Flowers on the Hills.

do. Chinensis splendid double mixed.

do. do. single mixed.

do. Heddewigi single and double finest mixed.

do. Imperialis double finest mixed.

Digitalis Purpurea (Fox glove).—Cold climate; finest mixed.

**Dimorphotheca** aurantiaca—A newly introduced splendid unnual for beds or pots. The colour of the flowers is deep orange yellow and the form of a daisy (As. 12).

Eschscholtzia (Californian Poppy).—A cold weather annual. Sow seeds like Poppy, but thinner. Foliage and flowers both ornamental and bright. Fine mixed single. (As. 6).

Forget-Me-Not Myosotis Alpestris.—SOW about end of rainy season.—SOIL damp garden soil.—WATER, daily. Flowers only in cold climate.

Fuchsia.—(For cold climate) hybrida double and single, splendid mixed As. 8.

Gaillardia Picta.—COLOUR, orange, red &c. HEIGHT one foot.—SEASON the whole year. TRANSPLANT 12 inches apart.—SOIL, rich and friable WATER, enough to keep soil moist. Good for cutting.

Gaillardia grandiflora-Splendid mixed.

do picta (Drummondii)

do. Lorenziana double mixed.

Gerbera Jamesoni, Transval daisy—A very showy flower of different shades of colour. It is perennial Per pkt. Re. 1.

Gladiolus Gandavensis, mixed colours. Produces roots like onions which last for years growing in size every year and producing in the rains bright coloured spikes. bearing a dozen or more orchid-like showy flowers.



Gloxinia

Gloxinia, splendid mixture of all classes (As. 8). The seed is small and the plant slow growing. Roots perennial.



Godetia

Godetia—COLOUR, crimson, lilac, rose and white,—HEIGHT 1 to 1½ feet. SOW after rains, and transplant into good soil 6 inches apart. Godetia, many sorts mixed.

do. Lady Albemarle (crimson).

do. Duchess of Albany (Satiny white.)

do. do. of Fife (white spotted carmine).



Globe Amaranth, Gomphrena Globosa

Gomphrena (Everlasting flowers) COLOUR, purple, white, pink, orange—SOW, beginning of monsoon till end.—SOIL, any soil not very rich—WATER, sufficient to keep soil moist.

Gomphrena globosa purple do. do. white



Gypsophila Paniculata.

Gomphrena rosea (pink). do. do. orange.

Gypsophila elegans pure white small decorative flowers.

Gypsophila paniculata pure white small decorative flowers. To be sown in the place where wanted to bloom in the rainy and cold season. Thin out to 6 in, apart,



Helianthus (Sun Flower—Colour cream, yellow, orange, red.—HEIGHT, 2 to 10 ft. SOW, June to November, in groups or at back of borders, if tall. The small kinds grow all the year round.

Helianthus Cucumerifolius.

Helianthus globosus fistulosus very double.

do. nanus fl. ple dwarf, double.

do. argyrophyllus (silver leaved).

do. macrophyllus giganteus (very large flowers).

do. cucumerifolius Diadem. cream white (good for cutting). ...

do. do. orion (crisped petals).

do. Japanese small (very useful for cutting).

do. Red (As. 8).

Helichrysum—Ever-lasting flowers—COLOUR, white, rose, yellow—HEIGHT 2 to 4 feet.—SOW, beginning to end of rainy season—SOIL, fair garden soil.

Helichrysum monstrosum, splendid double mixed.



Heliotropium,

do. mixed.

Heliotropium — (Heliotrope.) COLOUR purple shades with white and cream—HEIGHT 1 to 4 feet.—SOW, in and after rains. Transplant and protect from sun. Plants should be treated as perennials. In cool places they bear flowers all the year round.

Heliotropium dark vars, mixed.

do. Voltaireanum.

do. all sorts fine mixed.

do. large flowered.

Hollyhock—COLOUR, crimson, white, pink, &c.—HEIGHT five feet—SOW from beginning of rainy, to middle of cold season in beds of rich soil—Late sown plants bloom following year.

Hollyhock, finest double mixed (As. 6).

do. single do.

Impatiens Sultani and Holstii, hybrid, finest mixed. Zanzibar Balsam, Plants keep flowering the whole year. A good flower for decorating ferneries. This is one of the plants which flower best in shade. Sow hot or rainy season. Per pkt. Re. 1.

Ipomoea—COLOUR, dark blue, scarlet, red, yellow &c.— HEIGHT, 20 feet climbing, SEASON rainy and cold— SOW any time during rainy season where intended to

bloom.—SOIL deep and rich.

Ipomoea bona nox blue, ("Good night" flower).
do. grandiflora alba (mexicana), white.

do. quamoclit alba (white) and coccinea (red) Cypress Vine.

do. purpurea (morning glory), purple.

do. hederacea grandiflora superba, blue.

do. Leari (dark blue).



Larkspur— COLOUR, white, pink, purple and blue.—HEIGHT, 1 to 2 feet.—SOW, October, November and December.—SOIL any good soil. Early sowings fail.

Larkspur Dwarf double rocket mixed.

do. do. white (As. 6.)

do. Ranunculi-flowered, brick red.


do. Emperor, white (As. 6.)


do. do. Rose, blue and purple, (Separate colours). As. 6 per packet.

do. all colours mixed.



Linum grandiflorum rubrum (red) and roseum (pink).—A very showy cold season annual, with fine foliage and bright flowers. Should be massed in beds. Seed sown where wanted to bloom.





Lobelia erinus speciosa.

Lobelia erinus compacta.

Lobelia—COLOUR, blue, lilac, &c.—HEIGHT, 4 to 9 inches. -SEASON, cold to hot-SOW, if rainfall is light from middle of rainy season; if heavy, during cold season.— SOIL, rich friable.—WATER, enough to keep soil moist.

Lobelia erinus, (gracilis) blue and white. do. hybrida grandiflora, blue.

cardinalis, Queen Victoria, brilliant scarlet.

do. compacta, Crystal Palace.

pendula, (for hanging baskets), Per pkt. As. 12.

Lupinus, mixed annual tall sorts. A cold season annual having long sprays like larkspur, very useful for cutting. Grow like Larkspur.

Marigold—COLOUR, yellow and orange.—HEIGHT, 1 to 4 feet.—SEASON, rainy.—SOW, beginning to end of rains.—TRANSPLANT, when a few inches high.— WATER, keep soil moist.

Marigold, African, tall double, very large flowering.

do. do. '

dwarf mixed.

Prince of Orange and Lemon Queen. do.

do. French, mixed tall double and striped.


Mignonette—COLOUR, light green—HEIGHT, about a foot.— SOIL deep friable, containing abundance of lime .-SOW, after the heavy rains between August and November where wanted to bloom, in a partly shaded place.-TRANSPLANTING done gently, if necessary.

Mignonette, large flowering,

| Mignonette, fine sorts mixed do Machet,

do. Pyramidalis, gigantea do. Paris market.

excellent for pots.



Mimulus tigrinus

Mimulus.—COLOUR, yellow, maroon.—HEIGHT. 6 inches.—SEASON, cold.—SOW, in pots in a very fine soil, cover very slightly, stand seed pot in a saucer with water until germinated.—TRANSPLANT, 6 inches apart.

Mimulus cardinalis best sorts mixed | Mimulus tigrinus pardo, moschatus (musk plant). | dinus per pkt. As 8.



Mina Lobata.—A very fine annual climber. Flower shaded white, yellow and red, the sprays are very useful for table decorations.


per pkt. As. 12.

#### Mina Lobata.

Mirabilis Jalapa.—(Marvel of Peru 4 o'clock flower). COLOUR, white, yellow, crimson and spotted—HEIGHT, 2 to 4 feet. SEASON, rainy.—SOW, beginning of rainy season.—SOIL, rich and well-drained.—WATER, very profusely, one good watering daily, when not raining.

Mirabilis Jalapa Tom Thumb, striped leaved splendid mixed.

Nasturtium.—(Tropœlum)—COLOUR, orange, crimson, &c.—
HEIGHT, dwarf—1 foot. Tall 6 feet trailing or climbing—
SOW, rainy to cold season.—SOIL, ordinary garden.



Tropæolum (Nasturtium) Lobbianum.

Nasturtium, Lobbianum, finest mixed 8.

do. Majus, creeping, finest mixed.

do. do. atropurpureum (dark-crimson.)

do. do. coccineum (scarlet.)

do. do. Dunnett's orange (Sun-light.)

do. do. Pearl whitish, (Moonlight.)

do. do. Scarlet striped.

Nasturtium, Tom Thumb, finest mixture. Also following varieties.

do. Aurora (terra cotta).

do. Beauty, Yellow and Scarlet.

do. Coccineum (red.)

do. Crystal Palace Gem.

do. Empress of India.

do. Deep crimson dark leaved.

do. Golden Queen.

do. King Theodore.

do. Luteum (yellow)

do. Pearl white.

do. Prince Henry.

do, (yellow marbled searlet)


do. Purpureum (purple.)

do. Spotted.

do. Vesuvius red (Salmon.)







Nemophila.

Nemophila mixed, all sorts. A cold season annual. Very good for ribbon borders or small beds. Cultivate like Larkspurs.



Nicotiana.

Nicotiana (cold and hot season an-Tobacco plant nual). This is the bearing sweet scented flowers. SOW October, November, or earlier under shelter. Transplant 3 feet apart. Stake the flowering stalks.

affinis (white flowered Tobacco). sweet scented.

silvestris, long pure white flowers.


affinis hybrida, mixed colours.

do. Sanderae (reddish flowers.)

Nigella hispanica (Love in a mist). Nigella damascena (Devil-in-a-bush). A cold season annual. Nolana mixed (trailing annual)

Oenothera Drummondii (Evening Primrose) annual sorts mixed. Trailing habit Flowers pale yellow, open towards evening. SOW seeds at end of rains.

Transplant when six leaves have formed; flowers well in the hot season.



Pansy.

Pansy (Hearts ease)—CO-LOUR white, yellow, purple, &c.—HEIGHT, 4 to 8 inches.—SEASON, cold-SOW if rainfall is light, in middle of rainy season; if heavy, at end of rainy season-Water daily. Avoid afternoon sun. Can grow under shelter even during the rainy season.

Pansy maxima very fine mixture.

atropurpurea, dark purple. do: do.

atrosanguinea, dark red. do. do.

candidissima, white. do. do.

do. giant very large flowering Emperor do. William colour Corn flower blue-

do. Giant sorts mixd. do.

Per pkt. As. 8.

Pelargonium (Geranium) Zonal mixed fine.


do. silver foliaged (As. 8).

do, golden and bronze leaved (Re. 1).

do. scented leaved (As. 8).

Perilla Nankinensis—An annual with purple coloured and striped small sized leaves, like Coleus, 1-2 feet.

Petunia (Very good annual in places with moderate weather conditions)—COLOUR white. lilac, crimson, &c.—HEIGHT 1 to 3 feet, trailing.—SEAson end of rainy to hot.—SOW from middle of rainy to end of cold season, in pots, or seed Transplant when six heds. leaves are formed—WATER, keep the soil moist. Guard young plants against earth worms and slugs; the former eat roots, the latter the young leaves. Soot and ashes deter them.



#### Petunia Hybrida grandiflora.

Petunia hybrida inimitable striped and blotched.

do. do. pure white, very fine.

do. do. purpurea (purple).

do. do. illustris (rose carmine).

do. do. grandiflora single large flowering mixed (Re. 1),

do. superbissima dark rose.

do. do. fine mixed.

do. double large flowered (Rs. 1-8).

do. large flowering single fringed. do.

Phaseolus Caracalla (snail flower creeper) As. 8 per dozen see is.

A perennial creeper, bearing fragrant flowers, twisted in the form of a snail's shell. The plant can be trained on arches, grow like runner beans.



"Phlox Cuspidata" or Star Phlox.

Phlox Drummondif—CO-LOUR white, lilac, crimson, &c.—HEIGHT 6 inches to 1 foot—SEASON, nearly all the year round.
—SOW, if rainfall is light, from beginning of rainy to middle of cold season, if heavy, at the end of rainy season. WATER keep the soil moist. Transplant seedlings 6 to 9 inches apart.

C. Saberia T.

Phlox Drummondii finest mixture of all sorts.

do. do. grandiflora alba, pure white.

do. do. coccinea (red).

do. do. Kermesina (crimson).

do. do. Chamois rose.

do. do. atropurpurea deep purple.


do. do. stellata (centre white) finest mixed.

do. do. striata, striped vars, finest mixed.

do. do. cuspidata, Star Phlox, finest mixed (As. 6).

do. do. nana compacta (dwarf) extra fine mixed (As. 6).

do. perennis (Perennial finest mixed (Re. 1).



Poppy Dainty Lady.

42 14



Poppy Shirley.



Poppy.

Poppy.—COLOUR, white, rose, scarlet, &c.—HEIGHT, 9 to 15 inches.—SOW after rains in a loose rich soil.—Does not bear transplanting well. Too much water kills it.

Poppy, Iceland, double mixed.

do. do. single finest mixed.

do. Shirley, finest mixture.
do. Papaver orientale hybridum finest mixed.

do. Dainty Lady, rosy mauve.



Portulaca.—COLOUR, white, orange, crimson, &c.—HEIGHT, 3 to 6 inches creeping.—SEASON, end of rainy to end of hot.—SOW, from rainy to end of cold season, it rainfall light; if heavy, end of rainy season where intended to bloom or in pots.—SOIL should be well watered before sowing, then shaded till seeds are germinated. — WATER, keep soil moist. Flowers open from 8 to 5 by day.

Portulaca grandiflora single finest mixed.

do. do. splendid double mixed (Per pkt. As. 8).

Primula chinensis finest mixed. Cold season (Per pkt. Re. 1). For hill Stations.

Pyrethrum, Parthenifolium aureum (Golden feather). A hardy perennial As. 8.

Pyrethrum, Parthenifolium aureum laciniatum As. 8. do. roseum grandiflorum comet flowered As. 8.



#### Rhodanthe.

Rhodanthe single sorts fine mixed. Bears ever-lasting flower during the rainy and cold seasons.



Rudbeckia.

Rudbeckia.—Rainy season annual. Flowers, yellow and maroon good for cutting.

do. Amplexicaulis.

do. bicolour superba, large flowered.

do., Fulgida variabillis.

**Salpiglossis** finest mixed. Cold season annual with curious handsome flowers.



Salvia coccinea.

Salvia.—SOW in pots or a seedbed and transplant when the seedlings are 6 inches high about a foot apart.

Salvia argentea fine large silvery leaf.

do. coccinea fiery scarlet.

do. farinacea, bright, light

do. Horminum (Rose'clary) scarlet.

do. splendens scarlet, splendid

do. red varieties finest mixed

Saxifraga Sarmentosa (12 As.)—A good plant for hanging baskets.

Scabious (Pin cushion flower)—Dwar1 and semi-dwarf double mixed. Fine balls of flowers. Requires two seasons to blossom. (12 As. Per Pkt.)


Schizanthus, fine mixed. Butter-fly flower. Cold season annual. (Re. 1. Per Pkt.)

Senecio elegans (Jacobaea)—A hardy bedding annual, various colours.



Sensitive Plant

Sensitive Plant (Mimosa pudica)
Indian "Touch me not" The
leaves close up and droop down
with the slightest touch.



Smilax myrsiphyllum asparagoides. Produces glossy leaved trailers, useful for table decoration and bouquets.



Stock large flowering.

#### Stocks-Ten-Week COLOUR,


white, orange, scarlet, &c.— HEIGHT 6 inches to 1 foot. SEA-SON, cold weather.—SOW, after the heavy rains, preferably where wanted to bloom—WATER, carefully.

Stocks, German.

do. do. Dwarf bouquet Per pkt. As. 6.

do. do. large flowering dwarf finest mixture of all sorts. Per pkt. As. 6.


Streptocarpus achimeniflourus—A cold weather annual, with beautiful flowers like Achimenes.



Sweet Sultan.

Sultan, Sweet.—(Centaurea moschata)—Yellow-blue and white, separate colours or mixed. Flowers used for cutting. Cold season hardy annual. Better sown where intended to bloom after the rains.

Sweet Pea (Lathyrus odoratus)—SOW, from August to beginning of November, 6 inches apart, and in small groups or lines. Support on sticks. Or sow in pots 6 seeds, at equal distances, (and support on a thin trellis. Acglimatized seeds from hill stations do better in the plains.



Sweet Pea. Per pkt. As. 8.

Sweet Pea, Choicest, very rich.

do. Waved-standard varieties, splendid.

Avalanche, best white. do.

Brilliant Rose. do.

Floradale Fairy, exquisite primrose. do.

Fordhook Orange. do.

Gold Crest, rich salmon. do. Pimpernel, vivid scarlet. do.

Queen Alexandra Spencer, Vermillion Brilliant.

Sweet William (See Dianthus barbatus) extra fine mixed. Does well on the hills.

Tithonia Speciosa—A very hardy annual. Grows taller than Zinnias about 5 or 6 feet. Has orange red single flowers like Zinnias. Makes great effect in a distant border or clamp.


Torenia Fournieri—Nice plant for a bed or border.—SOW seeds in a pot and plant out when fit to handle. Enjoys a moist partly shady place. Makes a good edging. Rainy season. Torenia mixed.

Torenia Fournieri alba do. blue.

Bailloni yellow with do. brown.



Tropæolum Canariense Canary-creeper and service and the fed like creeping Nashurtiums,



Verbena hybrida.

Verbena Maminoth, grandifiora.

do. eitriodora, sweet scented.

do Drumentii.

do. Hybrida good quality mixed.

do. do. do. 1st quality

do. Candidissima pure white | Per pkt. As. 6. do.

do. do. Corrineo, scarlet.

do. do. firefiv. red.

do. pink shades very fine

do. Italian striped spl-edid.

de. Desert extra splendid mixed.

Verbens - COLOUR. white. purple. rose, &c-HEIGHT, 6 inches, creeping, SEASON, all the year round.-SOW any time during rains if rainfall is light, when rains are over, if heavy. -SOIL, rich and light with .. much leafmould as possible. often renewed.-WA-TER. keep the soil moist.

Verbesing .- a few tasts and the very round and earlier than the second flowers like Coreopsis. a frage to

Wiola Odorata seweet violet allea, white double As S
do. double blue do. As S
do. single time do. As S

Wallflower, quentid mixel. Flowers since it the possess.

Xeranthemum ever lasting houble white. Hairly and sold season annual. Sow seeds in pans, and plant out & necked apart. Good for bouquets.

Zinnia Elegans COLOUR, white, yellow, use in .—HEIGHT

10 inches to 4 feet.—SEASON tain; to middle of orbit.—

SOW any time during rains season, if minist is light:

near the end of rains, if heavy: transplant when a leaves

are formed.—WATER freely.

Zinnia elegans dere plane, tall deside.

do. scarlet
do. rosy pink.
do. crimson.
do. sulphur yellow.
do. white.

do. purple.

do. Lilliput fine, double mixed.

de. grandiflere rebusta, giant muxed As. 5.1

do. do. gold, striped scarlet (As. 6.)

do. Haageana hybrid fl. pleno (As. 12.)

do. crispa, curled and crested (As. 6

do. Picotti (Re. 1.)

#### **GRASS SEEDS**

Lawn grass seeds timported per line 2 d
Hariali or Dub grass 5 d
Lucerne, Poona grown (subject
to fluctuation of market Prices or application

## SEEDS OF USEFUL TREES AND SHRUBS As. 8 per Packet

(Yernacular)	(Botanical name)
Aboli (shrub Salmon Jowers)	Crossandra undulaefolia
Almorah Rohituka (tree) Ambadi, lal (shrub) Agasta (large flowers, tree)	F. Amoora Rohituka
Ambadi, lal (shrub)	Hibiscus sabdariffa
Agasta (large flowers, tree)	Sesbania grandiflora
	3 varieties.
Am (mara wooded tree)	. Terminana comencosa.
Asclepias (shrub beautiful flowers)	Asclepias curassavica
Asoka (ever green tree) Babool (gum arabica tree)	polyalthia longifolia
Babool (gum arabica tree)	Acacia Arabica
Bhendy tree (cart pole tree)	Thespesia populnea
Bhava (Indian Laburnum)	Cassia Fistula
Bor (Ber fruit tree)	
Buckool (ever green tree)	
Cassid (pink flowers, tree) vars	Cassia Marginata
do. (yellow flowers tree) vars Castor oil plant varieties	Disima amarana
Charden in the factor is a second of the factor is the factor in the factor is the factor in the factor in the factor is the factor in the fac	Dicinus communis
Chendufuli (the flower head is exa	
ly like a large drum stick, a shatree)	
Clerodendron (flowers brantiful)	··· Parkia biglandulosa
Cowit (wood apple) Curranja (Shade tree) Daisy tree (hears white flowers	···Clerodendron, varieties.
Currania (Shade tree)	Feronia elephantum
Daisy tree (bears white flowers	* Pongamia glabra
Christmas, like larke daisies	Montones O marieties
Dombeya (gives truses of flowers)	Dombova varieties
Gamphela (sprays of fragrant yello	· Donneya, varieues.
flowers, makes a fine back grou	ow nd
Con land	
	· · · · Galphimia glauca
Totropho Charm Co. C.	Holoptelea integrifolia
V and -1 () 1 1:6 1 ()	··· Jatropha, varieties ·· Bauhinia varieties
Kinhai (minati	· · Albizzia procera
Kinkari (sweet scented flowers,	a a
shrub)	Acacia Farnaciana
Kleinhovia (gives bunches of pinki	sh
nowers a small tree.)	···Kleinhovia hospita.
Mornand (nowers beautiful)	. Parlaria variation
remin (China box very sweet blo	OS-
soms)	··· Murraya exotica
Logwood (makes a close hedge)	Haematoxylon Campechi
	anum
Dibi dibi (pods yield tannin, a tree	) Caesalpinia coriaria

Mahogany (timber and avenue tree) Swietenia macrophylla and Mahogany.
Meetachinch (makes a thick hedge
which can be kept to any height) Ingadulcis
Meyena (fine large flowers, shrub) Meyenia, erecta
Myrtle (scented leaves and flowers, a
shrub) Myrtus communis
Nag Champa (fine bunches of yellow
flowers tall shrubs) Tecoma stans and Smithii
Pallas (Flame of the Forest, a tree) Butea frondosa
Pangara (Flame of the Forest, a tree) Erythrina indica and others
Parijataka (most sweet scented flow-
ers, a tree) Nyctanthus arbortristis
Peela Champa (sweet scenfed flow-
ers, a tree) Michelia Champaka
Peepul (the sacred tree) Figure religiosa
Peepul (the sacred tree) Ficus religiosa Putravanti (ever green tree) Putranjiva Roxburghii
Ritha (Soap nut tree) Sapindus trifoliatus
Sagwan (Teakwood tree) Tectona grandis
Shekta or Shevga (drum stick or In-
dian Horse Radish tree) Moringa pterigosperma
Shendri or Keshri Bixa orellana
Sirus (gigantic tree, blossoms very
strong scented) Albizzia Lebbek
Sissoo (Black wood tree) Dalbergia Sissoo
Tamarind (Imli tree) Tamarindus Indica
Tamhan (tree, mauve flowers)  Lagerstroemia flosreginae
Tarwad (Shrub, flowers yellow)  Cassia, varieties
Toon (avenue tree) Cedrela Toona
The state of the s

Hindu woman, a shrub)
SEEDS OF PLANTS SUITABLE FOR HEDGES

Tulsi (medicinal, sacred plant of the Ocimum, varieties.

			Seeds	
			Per oz.	Per lb.
(SA) - (S			As.	Rs.
Dodonaea viscosa		5. a. a.	4	3
Galphimia glauca			14	12
Duranta ellissii and plun	neria	1	3	2
Ingadulcis				2
Lawsonia alba	• • •		12	10
Sesbania aegyptiaca			3	2

Also, Agaves (commonly but wrongly called "Aloes"), Adhatoda (Adulsa), Euphorbia Tirucally (Milk bush), Jatropha curcas, Bamboo, Pomegranate, sour and wild lime or citron trees &c., which are propagated by seeds, cuttings or suckers, are recommended as good hedge plants.

and the transfer of the second second

skydyman ministra (and murva kas redució) genzedika Pontropolica in a

algerial for a solution in the solution of the

We want Office Americal Statement Statement of Committee and Committee a

the of the designation of the beam of the second of the se

New York and the second relies of resident and supported and

Marie Barrante succession of the College Hall Security

Colles of war of the Const. (nec) (which instituted the order of the collection of the Const. of the discrete discrete and others.

paradorna de la contra del la contra

Langli Levinera (care) animal) alsed

The management of the control of the

Survey (Bed loss) ... english grandlag ...

than Store tadich tree! - - : Moringa glerigosperma

The second secon

Ale in the second of the secon

consist processes to the contract of the contr


eminated the collection of the

and the second of the significant control of the second of

Separation of the property of the South

		A-min he are not a series
8 24		in a service of the s
	Proceedings of the	alah sidahara

Classic belief to the second of the second o



### ON SALE



#### PLANTS

Fruit trees.
Avenue trees.
Timber trees.
Fibre-yielding plants.
Other Economic plants.
Hedge plants.
Coffee plants.
Rose plants.
Foliage plants, in great variety.

Climbing plants of sorts.

Bulbs and tuberous-rooted plants.

Conservatory plants in great variety.

Palms and Cycads.

Ferns and Selaginellas.

Aquatic plants.

Edgings.

BOUQUETS, WREATHS, CROSSES AND CUT FLOWERS
GARDEN TOOLS.
INSECTICIDES.

AT THE

# EMPRESS BOTANICAL GARDENS

The Star Press, Poona.